
GLASSICE CHALK CRANK TWEAK CRYSTAL

From 2014-2017, there was a

38% INCREASE

in methamphetamine treatment admissions in

IOWA



BEFORE YOU RISK IT

GET THE FACTS

Meth causes brain changes that make it difficult to stop using.

THE

Meth use can cause a "crash" after the effects wear off. Meth KNOW use can cause long-lasting damage to the brain. Meth users RISKS who inject drugs and share needles are at risk for acquiring HIV and hepatitis.

1,713,000

persons in the U.S. aged 12 years or older reported using methamphetamine

High doses can elevate body temperature to dangerous, sometimes lethal, levels and cause convulsions and even cardiovascular collapse and death. Meth use may also cause extreme anorexia, memory loss and severe dental problems.

The use of methamphetamine during pregnancy carries significant risks. Research shows increased rates of premature delivery and placental abruption. Effects on babies include small size, lethargy, and heart and brain abnormalities.



FIND HELP NEAR YOU

Call (855) 582-8111. Text (855) 895-8398

You can contact Your Life Iowa 24/7 with guestions and concerns. Your call/text/chat is free and confidential.



1) Source: Iowa Department of Public Health (Disclaimer: Based on data reported by Iowa licensed treatment programs.) 2) Source: Iowa Department of Public Health (Disclaimer: Based on data reported by Iowa licensed treatment programs.) 3) Source: SAMHSA (Substance Abuse and Mental Health Service Administration) Tips for Teens 4) Source: DRUGS OF ABUSE | ADEA Resource Guide: 2017 Edition 5) Source: Annual Surveillance Report of Drug-Related Risks and Outcomes CDC National Center for Injury Prevention and Control, 2017 6) Source: National Institute on Drug Abuse

3